


Transverse Spin Results from PHENIX at RHIC

Feng Wei, for the PHENIX Collaboration
New Mexico State University

- ◆ Introduction
- ◆ PHENIX measurements and results
- ◆ Opportunities with new detectors

Single Transverse Spin Asymmetries



Theory Expectation (twist-2):

Small asymmetries at high energies

(Kane, Pumplin, Repko, PRL 41, 1689–1692 (1978))

$$A_N \propto \frac{m_q}{\sqrt{s}}$$

$A_N \sim \mathcal{O}(0.1\%)$ Theory


Experiment:

(E704, Fermi National Laboratory, 1991)

$$pp^{\uparrow} \rightarrow \pi + X$$






$$\sqrt{s} = 20 \text{ GeV}$$

$A_N \sim \mathcal{O}(10\%)$ Measured



How can we understand them?

Large Transverse Single Spin Asymmetry (SSA) in forward hadron production persists up to RHIC energy.



Non-Perturbative cross section

Perturbative cross section

7/17/2013




12th Asia Pacific
Physics Conference



3

Possible Origin of Large SSAs



Can initial and/or final state effects generate large transverse spin asymmetries?
($A_N \sim 10^{-1}$)

$$\frac{d^3 \sigma^\uparrow(pp^\uparrow \rightarrow \pi^+ X)}{dx_1 dx_2 dz} \propto \underbrace{q_i^\uparrow(x_1, k_{q,T}) \cdot G(x_2)}_{\text{Proton Structure}} \times \underbrace{\frac{d^3 \hat{\sigma}^\uparrow(q_i q_j \rightarrow q_k q_l)}{dx_1 dx_2}}_{\text{pQCD, small spin dependence}} \times \underbrace{FF_{q_{k,l}}(z, p_{h,T})}_{\text{fragmentation function}}$$

Mechanisms in QCD

TMD mechanism: GPM includes intrinsic transverse momentum

Possible Expansion:

$$A_N = \frac{\sigma^\uparrow - \sigma^\downarrow}{\sigma^\uparrow + \sigma^\downarrow} \propto f_{1T}^\perp \otimes D_1 + \delta q \otimes H_1^\perp + \dots$$

Sivers Function
(angular momentum)

↓

Sivers Effect

Transversity
(structure)

Collins Function
(Fragmentation)


↓

Collins Effect

Twist-3 Collinear factorization mechanism:


At high transverse momenta : two twist-3 correlation functions

1. Quark-gluon correlation function $T_{q,f}$
2. Two independent trigluon correlation functions $T_G^{(f)}, T_G^{(d)}$






The **R**elativistic **H**eavy **I**on **C**ollider
accelerator complex
at Brookhaven National Laboratory





RHIC as Polarized Proton Collider



PHENIX Detectors

2012



◆ Central Arm $|\eta| < 0.35$, $\Delta\phi = 2 \times \pi/2$

- Drift Chamber (DC)
- PbGl and PbSc (EMCal)
- Ring Imaging Cherenkov Detector (RICH)
- Pad Chambers (PC)
- Time Expansion Chamber (TEC)
- Silicon Vertex Detector (VTX)

◆ Muon Arms $1.2 < |\eta| < 2.4$, $\Delta\phi = 2\pi$

- Muon tracker (MuTr)
- Muon Identifier (MuID)
- RPC (Trig)
- Forward VTX (FVTX)

◆ Muon Piston Cal. (MPC) $3.1 < |\eta| < 3.9$

- Photons
- MPX-EX upgrade (2015)

◆ Global Detectors (Lumi, Trigger, local Pol.)

- BBC
- ZDC (neutron)

12th Asia Pacific
Physics Conference

PHENIX

Integrated Luminosity and Polarization

Year of RHIC Run	Energy [GeV]	Polarization [%]	Recorded L [pb ⁻¹]	FOM (P ² L) [nb ⁻¹]
2002	200	15	0.15	3.4
2005	200	47	0.16	35
2006	62.4	48	0.02	4.6
2006	200	50	2.7	700
2008	200	45	5.2	1100
2012	200	60	9.2	3300

PHENIX Measurements and Results

7/17/2013




12th Asia Pacific
Physics Conference



10


The Gluon Sivers measurement

- Can we separate the Sivers and Collins effects?



Heavy flavor production dominated by gluon-gluon fusion at RHIC energy

Gluon Fusion



- ◆ Eliminate the Collins' s effects since gluon has no transversity
- ◆ Sensitive to gluon Sivers function

Measurement of Heavy flavor decay muon

Muon Spectrometer:

● $1.2 < |\eta| < 2.4$

● Azimuthal: $\Delta\Phi = 2\pi$

$$A_N^{Phys} = \frac{A_N^{Incl} - r \cdot A_N^{BG}}{1 - r} \quad r = \frac{N^{BG}}{N^{Incl}}$$

Inclusive Muons

◆ Heavy Flavor decay muons


◆ Stopped hadrons → Distinguished background

→ Non-distinguished background

◆ Punch-through hadrons

◆ Hadron decay

muons



7/17/2013




12th Asia-Pacific
Physics Conference





12

A_N of heavy flavor decay muon



- ◆ Currently Measured A_N are consistent with zero.
- ◆ A larger 2012 data sample will increase sensitivity. The analysis is on progress.

◆ New installed FVTX detector will provide better rejection on hadron background in future




7/17/2013






12th A
Physics Conference



A_N of J/ψ




The results
are consistent
with zero too






A_N of Mid-rapidity π^0 and η

PRD 83, 032001 (2011)



Cross sections: pQCD in good agreement with RHIC data

7/17/2013





A_N : consistent with zero.

12th Asia Pacific
 Physics Conference



Forward A_N for MPC Clusters





Measurements from
MPC $3.1 < |\eta| < 3.9$

Cluster contribution

➤ Decay photon

➤ π^0

➤ Direct photon



7/17/2013

STATE


12th Asia Pacific
Physics Conference

PHENIX

Forward A_N Challenge: p_T Dependence

Valence Quarks' Sivers or Collins effects?


$x_F > 0.4$, Integrated Luminosity 33.0/pb, Polarization 0.60



- ◆ No sign of $1/p_T$ falloff yet.
 - Collins?
 - Twist-3 p_T dependent not trivial

- ◆ Much improved with MPC-EX (2015+)

Sub-process fractions p+p 200GeV



$$A_N \sim \frac{1}{Q} \quad @twist-3 \quad \text{Y. Koike, 2012}$$

$$A_N \sim O\left(\frac{M_N P_T S}{UT}\right) + O\left(\frac{M_N P_T}{-U}\right)$$

Interference Fragmentation

Interference Fragmentation Function (H_1):

Fragmentation of a transversely polarized quark q into two spin-less hadron h_1, h_2 carries an azimuthal dependence

$$d\sigma_{UT} = 2|P_{C\perp}||S_{BT}|\sin(\varphi_S - \varphi_R) \sum_{a,b,c,d} \int \frac{dx_a dx_b}{16\pi z_c} f_1^a(x_a) h_1^b(x_b) \frac{d\Delta\hat{\sigma}_{ab\rightarrow c\rightarrow d}}{d\hat{t}} H_{1,ot}^{qc}(\bar{z}_c, M_C^2)$$

$$A_{UT} = \frac{\sigma_{UT}}{\sigma_{UU}}$$

$$d\sigma_{UU} = 2|P_{C\perp}| \sum_{a,b,c,d} \int \frac{dx_a dx_b}{4\pi^2 z_c} f_1^a(x_a) f_2^b(x_b) \frac{d\hat{\sigma}_{ab\rightarrow cd}}{d\hat{t}} D_{1,00}(\bar{z}_c, M_C^2)$$

Transversity



IFF

Collins!

7/17/2


A. Bacchetta, M. Radici, PRD 70, 094032 (2004)

from polarization vector to scattering plane

$$\cos \varphi_S = \frac{(P_B \times P_C) \cdot (P_B \times S_B)}{|P_B \times P_C| \cdot |P_B \times S_B|}$$


from scattering plane to hadron plane

$$\cos \varphi_R = \frac{(P_C \times P_A) \cdot (P_C \times R_C)}{|P_C \times P_A| \cdot |P_C \times R_C|}$$





IFF Measurement

- ◆ PHENIX measurement in 2006 and 2008 are consistent with zero
- ◆ STAR has seen non-zero?
- ◆ Analysis of 2012 data is on-going,
- ◆ Move to more forward



7/17/2013

STATE



12th Asia Pacific
Physics Conference

PHENIX



PHENIX Opportunities with New Detectors

7/17/2013



12th Asia Pacific
Physics Conference





20

FVTX installed in 2012

4 layers of FVTX end-caps cover $1.2 < |\eta| < 2.4$ with resolution $\sim 100 \mu\text{m}$.

- ◆ Precise Charm/Beauty, W/Z Measurements
- ◆ Drell-Yan, J/ψ ... via dimuons



7/17/2013







12th Asia Pacific
Physics Conference

PHENIX

21

Projected Open Charm A_N with FVTX

- ◆ Significant rejection of hadron background
- ◆ Limited power on D/B separation
- ◆ Transverse spin data expected from 2015



7/17/2013




12th Asia Pacific
Physics Conference



22

Drell-Yan Process

- ◆ No fragmentation (no Collins effect folded)
- ◆ TMD factorization is valid
- ◆ The Sivers function in DY is opposite to that in DIS which can test current SSA QCD mechanisms (TMD & Twist-3)
- ◆ A_{TT} in Drell-Yan is sensitive to transversity!




$$p + p \rightarrow l^+ + l^- + X$$



$$l + p \rightarrow l + \pi + X$$

$$\Delta f_{1T}^\perp(x, \vec{k}_T^2)|_{\text{Drell-Yan}} = -\Delta f_{1T}^\perp(x, \vec{k}_T^2)|_{\text{SIDIS}}$$

Drell-Yan Projection



RHIC 1-year running projection



7/17/2013


STATE

12th Asia Pacific
Physics Conference

PHENIX

24

Coming soon: MPC-EX (2015+)





A combined charged particle tracker and EM pre-shower detector – dual gain readout allows sensitivity to MIPs and full energy EM showers.

$$3.1 < |\eta| < 3.9$$

- π^0 rejection \rightarrow direct photons
- π^0 reconstruction out to $>80\text{ GeV}$

7/17/2013




12th Asia Pacific
Physics Conference




Polarized p+A at RHIC(2015+)

- ◆ Large transverse spin asymmetry A_N at forward rapidity - a large analyzing power at large x_1
- ◆ Gluon saturation/CGC probed at forward rapidity in p+A - small x_2 in A



projectile: $x_1 \sim \frac{p_{\perp}}{s} e^{+y} \sim 1$ valence


target: $x_2 \sim \frac{p_{\perp}}{s} e^{-y} \ll 1$ gluon



$$\Delta\sigma_{\text{forward}} \sim \Delta f(x_1) \otimes g(x_2); \quad x_1 \gg x_2$$

Forward Pion Single-Spin Asymmetry

$$\frac{A_N^{pA \rightarrow h}}{A_N^{pp \rightarrow h}} \bigg|_{P_{h\perp}^2 \ll Q_s^2} \approx \frac{Q_{sp}^2}{Q_{sA}^2} e^{\frac{P_{h\perp}^2}{Q_{sp}^2} \frac{\sigma^2}{4}}$$



12th Asia Pacific Physics Conference

Kang, Yuan (2011)

PHENIX

Summary and Outlook

- ◆ **PHENIX has measured A_N in heavy flavor production**
 - Statistics is still limited and 2012 data will be included
 - Previous A_N of J/ψ was published and consistent with 2012 data
- ◆ **A_N of π^0 and η at mid- and forward rapidity was measured**
 - Current results prepare to be published
 - Expected much improved measurements with MPC-ex
- ◆ **A_N from Interference Fragmentation was measured**
 - Expect 2012 result soon with more forward measurement
- ◆ **Forward measurements with FVTX**
 - Expect significant background projection in heavy flavor production
 - Drell-Yan measurement becomes possible
- ◆ **More opportunities with polarized p+A experiments in future**

Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil

Institute of Physics, Academia Sinica, Taipei 11529, Taiwan

China Institute of Atomic Energy (CIAE), Beijing, People's Republic of China

Peking University, Beijing, People's Republic of China

Charles University, Ovocnyth 5, Praha 1, 116 36, Prague, Czech Republic

Czech Technical University, Zikova 4, 166 36 Prague 6, Czech Republic

Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2,

182 21 Prague 8, Czech Republic

Helsinki Institute of Physics and University of Jyväskylä, P.O.Box 35, FI-40014 Jyväskylä, Finland

Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France

Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay,

F-91128, Palaiseau, France

Laboratoire de Physique Corpusculaire (LPC), Université Blaise Pascal, CNRS-IN2P3,

Clermont-Fd, 63177 Aubiere Cedex, France

IPN-Orsay, Université Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France

Debrecen University, H-4010 Debrecen, Egyetem tér 1, Hungary

ELTE, Eötvös Loránd University, H - 1117 Budapest, Pázmány P. s. 1/A, Hungary

KFKI Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences (MTA KFKI RMKI),

H-1525 Budapest 114, POBox 49, Budapest, Hungary

Department of Physics, Banaras Hindu University, Varanasi 221005, India

Bhabha Atomic Research Centre, Bombay 400 085, India

Weizmann Institute, Rehovot 76100, Israel

Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo,

Tokyo 113-0033, Japan

Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan

KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan

Kyoto University, Kyoto 606-8502, Japan

Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan

RIKEN, The Institute of Physical and Chemical Research, Wako, Saitama 351-0198, Japan

Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan

Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan

Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

Chonbuk National University, Jeonju, Korea

Ewha Womans University, Seoul 120-750, Korea

Hanyang University, Seoul 133-792, Korea

KAERI, Cyclotron Application Laboratory, Seoul, South Korea

Korea University, Seoul, 136-701, Korea

Myongji University, Yongin, Kyonggido 449-728, Korea

Department of Physics and Astronomy, Seoul National University, Seoul, South Korea

Yonsei University, IPAP, Seoul 120-749, Korea

IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics,

Protvino, 142281, Russia

INR_RAS, Institute for Nuclear Research of the Russian Academy of Sciences, prospekt 60-letiya Oktyabrya 7a,

Moscow 117312, Russia

Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia

Russian Research Center "Kurchatov Institute", Moscow, Russia

PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia

Saint Petersburg State Polytechnic University, St. Petersburg, Russia

Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Vorob'yev Gory,

Moscow 119992, Russia

Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden

PH ENIX

13 Countries; 70 Institutions



Thanks



7/17/2013

STATE

Physics Conference

PH ENIX

28

Back up

7/17/2013



12th Asia Pacific
Physics Conference



29


Spin in transversely polarized proton

- Quark transversity
 - Know much better about quark transversity than before.
- Gluon transversity
 - No transversely polarized gluon.
- Parton orbital angular momentum
 - The Sivers function could be related to orbital angular momentum.
 - Quark Sivers function is constrained OK.
 - Gluon Sivers function is not well known.

7/17/2013



12th Asia-Pacific
Physics Conference



M. Anselmino
et. al.

Nucl. Phys.
Proc. Suppl.
191, 98
(2009).

Eur. Phys. J.
A39:89–
100, 2009


PHENIX

Possible Mechanisms for A_N

- Quarks' Sivers and Collins TSSA observed in SIDIS
- Gluons' Sivers not constrained in SIDIS @LO

Sivers mechanism: Correlation between nucleon spin and parton k_T

Phys Rev D41 (1990) 83; 43 (1991) 261



Forward focus:

- Valence quark Sivers via pion
- Gluon Sivers via heavy quarks
- Drell-Yan Sivers sign change


Orbital Angular Momentum?

7/17/2013



Collins mechanism: Transversity (quark polarization) * Spin-dependence in the jet fragmentation

Nucl Phys B396 (1993) 161



Requires full jet measurements
- forward s/ePHENIX upgrade

12th Asia Pacific
Physics Conference




31

Semi-classical Interpretation

Two Possibilities:

1. Nuclear Shadowing
(angular momentum)







$u(x, \mathbf{b}_\perp)$

$u_X(x, \mathbf{b}_\perp)$


$d(x, \mathbf{b}_\perp)$

$d_X(x, \mathbf{b}_\perp)$



2. Impact Parameter Space Changes with polarization

- ◆ Attractive rescattering of hit quark by gluon creates transverse momentum
- ◆ M.Burkardt [hep-ph0309269] – impact parameter formalism
- ◆ Orbital angular momentum at finite impact parameter
- ◆ Observed and true x differ
- ◆ Observable left/right asymmetry



Heavy Quark TSSA at RHIC

Twist-3 tri-gluon correlation functions

$$P_h^0 \frac{d\sigma^{3\text{gluon}}}{d^3P_h} \simeq \frac{\alpha_s^2 M_N \pi}{S} \epsilon^{P_h p n S_\perp} \sum_{f=c\bar{c}} \int \frac{dx'}{x'} G(x') \int \frac{dz}{z^3} D_a(z) \int \frac{dx}{x} \delta(\tilde{s} + \tilde{t} + \tilde{u}) \frac{1}{\tilde{u}} \times \left[\delta_f \left(\frac{d}{dx} O(x) - \frac{2O(x)}{x} \right) \hat{\sigma}^{O1} + \left(\frac{d}{dx} N(x) - \frac{2N(x)}{x} \right) \hat{\sigma}^{N1} \right].$$

where $O(x) \equiv O(x, x) + O(x, 0)$, $N(x) \equiv N(x, x) - N(x, 0)$.

$\delta_f = +1(c); -1(\bar{c})$

?




$$A_N(D) \neq A_N(\bar{D})$$

Model 1:

$$O(x) = 0.004xG(x)$$

Koike *et. al.* (2011)

Kang, Qiu, Vogelsang, Yuan (2008)



7/17/2013




Physics Conference



33

A_N for Di-hadron



- Sivers asymmetry ($q_T \perp$)
- No asymmetry expected for $q_T \parallel$
- Improved statistics for 2008 data set!


TMD is not valid for back to back di-hadron process in p+p collision

Similar analysis possible in different combinations of rapidity

η_{\min}	-3.7	-2.4	-0.35	1.2	3.1
η_{\max}	-3.1	-1.2	+0.35	2.4	3.9

Forward A_N for Charged Hadrons

A_N for hadrons



- Measured at Muon Arms
 $1.2 < |\eta| < 2.4$
- No PID - Unidentified charge hadrons

